



Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com



Research letter

Evacuation is a risk factor for diabetes development among evacuees of the Great East Japan earthquake: A 4-year follow-up of the Fukushima Health Management Survey

Introduction

The Great East Japan earthquake struck on 11 March 2011. After the earthquake, a massive tsunami hit the Tokyo Electric Power company's Fukushima Daiichi nuclear power plant in Fukushima prefecture, causing the release of radiation. The Fukushima Daiichi nuclear disaster forced the evacuation of several towns, which led to lifestyle changes and anxiety over radiation among the evacuees. For this reason, immediately after the disaster, the Fukushima Health Management Survey (FHMS) was started to investigate the effects of long-term exposure to low-dose radiation [1].

Based on previous data from a comprehensive health check-up with a 1.6-year follow-up, it had been reported that evacuation was a cause of body weight gain [2], diabetes [3], hypo-high-density lipoprotein (hypo-HDL) cholesterolaemia [4], hypertension [5] and metabolic syndrome [6]. These results suggested that evacuation might be a risk factor for the development of various disorders.

In the previous study [3], after adjusting for age, gender, body mass index (BMI), smoking status, systolic blood pressure, HDL cholesterol (HDL-C), alanine aminotransferase (ALT) and γ -glutamyl transpeptidase (γ -GT) levels using the Cox proportional hazards model, it was demonstrated that evacuation was associated with a significantly higher incidence of diabetes. As it was unclear why this was so, it was assumed that the reason might be the associated change in lifestyle with evacuation. Lifestyle change is a well-known cause of obesity that, in turn, can lead to an increase in incidence of diabetes.

Therefore, the present study concentrated more on lifestyle factors at baseline to determine whether they could confound the relationship between incident diabetes and evacuation, and also followed people over a longer time span of 4 years. Data from the aftermath of the disaster were used as baseline.

Methods

Study population and design

The FHMS was carried out by Fukushima Medical University. The participants were Japanese adults living near the Fukushima Daiichi nuclear power plant in Fukushima Prefecture who were residents of Tamura city, Minamisoma city, Kawamata town, Hirono town, Naraha town, Tomioka town, Kawauchi village, Okuma town, Futaba town, Namie town, Katsurao village, Iitate

village and Date city. According to census data from 2010, the combined population of these communities was 278,286 individuals. In 2008, the Japanese government had started an annual health-check programme, The Specific Health Check and Guidance System, which focused on detecting metabolic syndromes in adults aged 40–74 years and was run by national healthcare insurers. The dates of annual health check-ups from 2011 as a part of the comprehensive health checks in the FHMS were followed. A previous report details the methodology of these health checks and the FHMS [1]. Based on national census data, the target population of adults aged 40–74 years living in the above-mentioned 13 municipalities was 125,987 in 2010, a year before the disaster. Also obtained were data for 21,354 adults who had undergone annual health check-ups in 2011 after the disaster (baseline) as well as at least one check-up between 2012 and 2014; they represented around 17% of the population according to the 2010 census. To conduct a longitudinal analysis, 3001 participants who had not received an annual check-up between 2012 and 2014 were excluded. Also excluded were 1989 participants who had been diagnosed with diabetes in 2011 check-ups, and 2877 participants who were missing data on, for example, physical activity (2798 participants), weight changes from age 20 years (46 participants), weight changes within one year (24 participants) and hours of restorative sleep (nine participants) as baseline. Finally, 13,487 participants who had undergone follow-up examinations were included in the analysis (Fig. S1; see supplementary data associated with this article online). For those who underwent check-ups more than once during the survey period, the date on which diabetes was identified, or the latest date in subjects without diabetes, was recorded.

Ethics and consent

Informed consent was obtained from all subjects and community representatives as required to conduct an epidemiological study as per the guidelines of the Council for International Organizations of Medical Science [7]. Also, the study protocol was reviewed and approved by the Ethics Committee of Fukushima Medical University (#1916), and conducted in accordance with the approved guidelines.

Measurements

All measurements used in this analysis are described in detail in the Appendix (see supplementary data associated with this article online).

Statistical analysis

Participants were divided into two groups, based on their residence status after the Great East Japan Earthquake, as evacuees

<https://doi.org/10.1016/j.diabet.2017.09.005>
1262-3636/© 2017 Elsevier Masson SAS. All rights reserved.

($n = 4235$) and non-evacuees ($n = 9252$). Baseline characteristics of the evacuees and non-evacuees who underwent follow-up health examinations were compared using chi-square or Wilcoxon rank-sum tests. To evaluate the impact of evacuation on incidence of diabetes, hazard ratios (HRs) of new-onset diabetes and 95% confidence intervals (CIs) for evacuation were calculated using multivariable-adjusted HRs after adjusting for other potential confounding factors [age, gender, BMI scores, current smoker, excess alcohol consumption, weight change (≥ 10 kg) from age 20 years, weight change (≥ 3 kg) within 1 year, sleep quality and physical activity]. SAS version 9.3 software (SAS Institute, Cary, NC, USA) was used for all analyses. All probability values for statistical tests were two-tailed, and P values < 0.05 were considered statistically significant.

Results

Baseline characteristics of participants who had follow-up health checks after the disaster

Among all 18,353 participants (including 1989 patients with diabetes), the prevalence of diabetes at baseline was estimated to be 10.8% (Supplementary data, Fig. S1). The study followed 13,487 participants (43.5% male) without diabetes using data from annual check-ups in 2010–2011 and those who had undergone at least one other annual check-up in 2012–2014 (Supplementary data, Fig. S1). Table 1 presents the clinical characteristics of the two groups (evacuees and non-evacuees) at baseline. First, when risk factors for the development of diabetes were compared, the evacuees were significantly younger and more often female than non-evacuees. The evacuees also had a significantly greater prevalence of obesity, dyslipidaemia, weight change (≥ 10 kg) from age 20, weight change (≥ 3 kg) within one year, adequate sleep and smoking habit than non-evacuees. However, no significant differences were observed in HbA1c levels, hypertension or physical activity between the two groups.

Table 1
Baseline characteristics of evacuees and non-evacuees who underwent follow-up health examinations after the Great East Japan Earthquake.

	Non-evacuees	Evacuees	P^a
n	9252	4235	
Gender (% male)	44.2%	41.9%	< 0.001
Age (years)	64.3 ± 7.8	63.1 ± 8.9	< 0.001
HbA1c (%)	5.3 ± 0.3	5.3 ± 0.3	0.071
HbA1c (mmol L ⁻¹)	34.4 ± 3.3	34.4 ± 3.3	0.071
Obesity (%) ^b	26.7	36.0	< 0.001
Hypertension (%) ^c	52.7	51.9	0.374
Dyslipidaemia (%) ^d	50.0	56.0	< 0.001
Weight change (≥ 10 kg) from age 20 (%)	31.3	39.6	< 0.001
Weight change (≥ 3 kg) within 1 year (%)	19.2	40.1	< 0.001
Regular exercise (%)	34.5	34.3	0.847
Adequate sleep (%)	75.1	64.0	< 0.001
Current smoker (%)	12.2	15.1	< 0.001
Alcohol consumption (%)			
Non-drinker	29.5	33.8	< 0.001
Light drinker	51.9	48.2	
Moderate/heavy drinker ^e	18.6	18.0	

Data are presented as means \pm standard deviation (SD) and as percentages for categorical variables.

^a Evacuees vs non-evacuees after earthquake by chi-square or Wilcoxon rank-sum test.

^b Body mass index ≥ 25 kg/m².

^c Systolic blood pressure ≥ 140 mmHg, diastolic blood pressure ≥ 90 mmHg or self-reported use of blood-pressure-lowering drugs.

^d LDL-C ≥ 140 mg dL⁻¹, triglycerides ≥ 150 mg dL⁻¹, HDL-C < 40 mg dL⁻¹, or self-reported use of cholesterol-lowering drugs.

^e Ethanol intake ≥ 44 g day⁻¹.

Incidence of diabetes

On investigating the incidence of diabetes, the cumulative incidence was 4.54% (612 participants) over a mean follow-up period of 2.67 years. The total incidence of diabetes was 17.0/1000 person-years, while that of the evacuees was 1.61-fold higher (23.2/1000 person-years) compared with non-evacuees (14.4/1000 person-years). Significant differences were also observed in baseline characteristics between the two groups, prompting the construction of Cox proportional hazards models, which found that evacuation was indeed a significant risk factor for the development of diabetes compared with non-evacuation, with a crude HR of 1.73 (95% CI: 1.47–2.04). The age- and gender-adjusted HR for evacuation was 1.78 (95% CI: 1.51–2.09) whereas, after full adjustment, the multivariate HR for evacuation was 1.51 (95% CI: 1.28–1.79).

Discussion

Our team had previously demonstrated that, after adjusting for age, gender, BMI, smoking status, and systolic blood pressure, HDL-C, ALT and γ -GT levels [3], evacuation was significantly associated with an increased incidence of diabetes. However, there were several limitations in that previous analysis. First, the results had been obtained from a short-term follow-up of only 1.6 years. Second, that analysis was not adjusted for lifestyle factors, such as weight change, physical activity and sleep quality. Therefore, the present study investigated the effects of prolonged evacuation on the incidence of diabetes after adjusting for these and other lifestyle factors. The results have revealed that evacuation after the disaster was an independent risk factor for the development of diabetes, even after adjusting for various lifestyle factors.

The present data have also shown that metabolic factors, including obesity and dyslipidaemia, adversely affected evacuees. In addition, ratios of weight change from age 20, sleep deprivation and current smoking were significantly higher in evacuees than in non-evacuees. Thus, the disaster was more likely to have negative effects on metabolic laboratory test results in evacuees compared with non-evacuees. These findings suggest that chronic metabolic health problems such as obesity, type 2 diabetes, hypertension and dyslipidaemia should be carefully monitored and treated following a disaster, especially among evacuees. Living as an evacuee under unfavourable conditions increases stress in terms of privacy, availability of food, duty assignments, income, jobs and health [8]. Particularly in diabetes patients, mental stress can aggravate diabetes control [8,9].

Although worsening of glycaemic control after a disaster may be affected by a number of factors, such as changes in diet, reduction in exercise and more psychological stress, the association between evacuation and increased diabetes incidence remains unclear. In a previous report of mental health care, the FHMS revealed that evacuees who believed that radiation exposure causes negative health effects were significantly more likely to be psychologically distressed [10]. In fact, psychological distress was significantly more prevalent among residents of evacuation zones even after adjusting for other significant risk factors, such as age, gender, living arrangements, experiencing the nuclear-power-plant accident, loss of a family member, becoming unemployed and a history of mental illness. Moreover, psychological distress in each evacuation zone was positively associated with radiation levels in the evacuees' environment [11].

These reports suggest that fear of radiation risk as well as socioeconomic factors contribute to psychological distress among evacuees. Although not conducted under natural-disaster conditions, a prospective population-based study in Finland showed that

high levels of psychological distress at baseline predict the development of metabolic syndrome, independent of age, gender, marital status, educational attainment, and baseline health behaviours such as smoking, alcohol intake and leisure-time physical activity [12]. Likewise, the FHMS found that evacuation was significantly associated with an increased incidence of metabolic syndrome [6]. Moreover, the survey also revealed that evacuation is a significant risk factor for overweight [2] and hypertension [5]. Thus, evacuees tend to be more psychologically distressed and are at greater risk of developing metabolic disorders than non-evacuees.

In conclusion, the findings of the present study suggest that prolonged evacuation is a risk factor for the development of diabetes following a disaster. Therefore, it is important to follow-up evacuees and to recommend lifestyle changes where necessary.

Disclosure of interest

The authors declare that they have no competing interest.

Acknowledgments

This survey was supported by the National Health Fund for children and adults affected by the nuclear incident. The findings and conclusions of this article are the sole responsibility of the authors, and do not represent the official views of the Fukushima Prefecture government.

Appendix A. Supplementary data

Supplementary data (Fig. S1 and Appendix) associated with this article can be found, in the online version, at <http://www.sciencedirect.com> at <http://dx.doi.org/10.1016/j.diabet.2017.09.005>.

References

- [1] Yasumura S, Hosoya M, Yamashita S, Kamiya K, Abe M, Akashi M, et al. Study protocol for the Fukushima Health Management Survey. *J Epidemiol* 2012;22:375–83.
- [2] Ohira T, Hosoya M, Yasumura S, Satoh H, Suzuki H, Sakai A, et al. Effect of evacuation on body weight after the Great East Japan Earthquake. *Am J Prev Med* 2016;50:553–60.
- [3] Satoh H, Ohira T, Hosoya M, Sakai A, Watanabe T, Ohtsuru A, et al. Evacuation after the Fukushima Daiichi nuclear power plant accident is a cause of diabetes: results from the Fukushima Health Management Survey. *J Diabetes Res* 2015;2015:627390.
- [4] Satoh H, Ohira T, Nagai M, Hosoya M, Sakai A, Watanabe T, et al. Hypo-high-density lipoprotein cholesterolemia caused by evacuation after the Fukushima Daiichi nuclear power plant accident: results from the Fukushima Health Management Survey. *Intern Med* 2016;55:1967–76.
- [5] Ohira T, Hosoya M, Yasumura S, Satoh H, Suzuki H, Sakai A, et al. Evacuation and risk of hypertension after the Great East Japan earthquake: the Fukushima Health Management Survey. *Hypertension* 2016;68:558–64.
- [6] Hashimoto S, Nagai M, Fukuma S, Ohira T, Hosoya M, Yasumura S, et al. Influence of post-disaster evacuation on incidence of metabolic syndrome. *J Atheroscler Thromb* 2017;24:327–37.
- [7] American Society of Law & Medicine. International guidelines for ethical review of epidemiological studies *Law Medicine & Health care: a publication of the American Society of Law & Medicine*, 19. 1991;p. 247–58.
- [8] Inui A, Kitaoka H, Majima M, Takamiya S, Uemoto M, Yonenaga C, et al. Effect of the Kobe earthquake on stress and glycemic control in patients with diabetes mellitus. *Arc Intern Med* 1998;158:274–8.
- [9] Hanson CL, Henggeler SW, Burghen GA. Model of associations between psychosocial variables and health-outcome measures of adolescents with IDDM. *Diabetes Care* 1987;10:752–8.
- [10] Suzuki Y, Yabe H, Yasumura S, Ohira T, Niwa S, Ohtsuru A, et al. Psychological distress and the perception of radiation risks: the Fukushima health management survey. *Bull World Health Organ* 2015;93:598–605.
- [11] Kunii Y, Suzuki Y, Shiga T, Yabe H, Yasumura S, Maeda M, et al. Severe psychological distress of evacuees in evacuation zone caused by the Fukushima Daiichi nuclear power plant accident: the Fukushima Health Management Survey. *PLoS One* 2016;11:e0158821.
- [12] Puustinen PJ, Koponen H, Kautiainen H, Mantyselka P, Vanhala M. Psychological distress predicts the development of the metabolic syndrome: a prospective population-based study. *Psychosom Med* 2011;73:158–65.

H. Satoh^{a,b,*}, T. Ohira^{b,c}, M. Nagai^{b,c}, M. Hosoya^{b,d}, A. Sakai^{b,e}, S. Yasumura^{b,f}, A. Ohtsuru^{b,g}, Y. Kawasaki^{b,d}, H. Suzuki^{b,h}, A. Takahashi^{b,i}, Y. Sugiura^{b,j}, H. Shishido^{b,k}, Y. Hayashi^{b,l}, H. Takahashi^{b,m}, G. Kobashi^{b,n}, K. Ozasa^{b,o}, S. Hashimoto^b, H. Ohto^b, M. Abe^b, K. Kamiya^{b,p}, Fukushima Health Management Survey Group

^aDepartment of Metabolism and Endocrinology, Juntendo University, 2-1-1 Hongo, Bunkyo-ku, 113-8421 Tokyo, Japan

^bRadiation Medical Science Center for the Fukushima Health Management Survey, Fukushima Medical University, Fukushima, Japan

^cDepartment of Epidemiology, Fukushima Medical University, Fukushima, Japan

^dDepartment of Pediatrics, Fukushima Medical University, Fukushima, Japan

^eDepartment of Radiation Life Sciences, Fukushima Medical University, Fukushima, Japan

^fDepartment of Public Health, Fukushima Medical University, Fukushima, Japan

^gDepartment of Radiation Health Management, Fukushima Medical University, Fukushima, Japan

^hDepartment of Cardiology and Hematology, Fukushima Medical University, Fukushima, Japan

ⁱDepartment of Gastroenterology and Rheumatology, Fukushima Medical University, Fukushima, Japan

^jDepartment of Neurology, Fukushima Medical University, Fukushima, Japan

^kDepartment of Orthopedic Surgery, Fukushima Medical University, Fukushima, Japan

^lDepartment of Nephrology, and Hypertension, Fukushima Medical University, Fukushima, Japan

^mInformation Management and Statistics Office, Radiation Medical Science Center for the Fukushima Health Management Survey, Fukushima Medical University, Fukushima, Japan

ⁿDepartment of Public Health, Dokkyo Medical University, Tochigi, Japan

^oDepartment of Epidemiology, Radiation Effects Research Foundation, Hiroshima, Japan

^pResearch Institute for Radiation Biology and Medicine, Hiroshima University, Hiroshima, Japan

*Corresponding author. Department of Metabolism and Endocrinology, Juntendo University 2-1-1 Hongo, Bunkyo-ku, Tokyo 113-8421, Japan

E-mail address: hk-sato@juntendo.ac.jp (H. Satoh).

Received 7 July 2017

Received in revised form 20 September 2017

Accepted 25 September 2017

Available online xxx